

What is Ganirelix Gedeon Richter and what is it used for?

Ganirelix Gedeon Richter is used to prevent premature LH surge that can cause premature ovulation in women undergoing assisted reproduction.

Ganirelix Gedeon Richter pre-filled syringe with fixed needle should be injected into the fat layer directly under the skin, as prescribed by your doctor or nurse. Do not mix Ganirelix Gedeon Richter with other medicines. Take your injection at the same time as you take your follicle-stimulating hormone.




How to use Ganirelix Gedeon Richter

Always take Ganirelix Gedeon Richter as prescribed by your doctor or nurse.

Prepare the injection

1

Read the leaflet carefully. Wash your hands and take a pre-filled syringe out of the package. Hold the syringe in the middle. Check that the solution in the syringe is clear and free of particles. Small air bubbles may be visible but these do not need to be removed. Hold the syringe horizontally with the needle pointing away from you and pull off the needle cover. Note: *Do not twist off the needle cover!*



Inject the dose

2

Inject the liquid (normally in your tummy) subcutaneously i.e into the fatty layer under the skin according to instructions from your doctor or nurse. Lightly pinch the skin of the injection area and insert the needle. Slowly and firmly push the plunger all the way down until the syringe is empty. Wait 5 seconds, then release the skin fold and withdraw the needle.



After the injection: Do not replace the needle cover to avoid needle stick injury.

Discard the used syringe safely in the puncture-resistant container and return to the pharmacy for a correct disposal. Ask your pharmacist how to dispose of medicines you no longer use.



How to store Ganirelix Gedeon Richter?

Do not freeze.

Store at room temperature in the original packaging.

Light sensitive.

If you forget to use Ganirelix Gedeon Richter

If you miss a dose, take it as soon as possible.

Do not take a double dose to compensate for a missed dose.

Contact your doctor or nurse if you are unsure.

Read the leaflet carefully
before you use
Ganirelix Gedeon Richter

Ganirelix Gedeon Richter (ganirelix), 0,25 mg/0,5 ml solution for injection in pre-filled syringe. Ganirelix Gedeon Richter is used to prevent the premature luteinising hormone surge that might cause premature release of egg cells in women undergoing assisted reproduction techniques.

Warnings and precautions: Do not use Ganirelix Gedeon Richter if you are allergic to ganirelix or any of the other ingredients of this medicine. If you have an active allergic condition or an allergic reaction, stop using Ganirelix Gedeon Richter and seek immediate medical assistance. Do not use the medicine if you are hypersensitive to gonadotropin-releasing hormone (GnRH) or a GnRH analogue or have a moderate or severe kidney or liver disease. You should not use Ganirelix Gedeon Richter if you are pregnant or breast-feeding. The most common side effects are local skin reactions at the site of injection. Less common side effects are headache, nausea and feeling generally unwell (malaise). **Carefully read the package leaflet before you use Ganirelix Gedeon Richter. It contains information that is important for you. Ask your doctor or nurse if you have any further questions and always follow their instructions. For more information, see the package leaflet or www.fass.se.**

Contact: Gedeon Richter Nordics AB, Barnhusgatan 22, 111 23 Stockholm.
Tel: 08-611 24 00. **Medical information:** medinfo.se@gedeonrichter.eu,
tel: 08-611 24 22. Abbreviated information is based on the package leaflet dated 2022-07-15.



GEDEON RICHTER

Barnhusgatan 22, 111 23 Stockholm, Sweden.

