


Patient instructions

Menopur[®]


600 IU / 1200 IU



Menopur[®]
menotropin

Be sure you have everything needed at hand before you begin:

- 1 Vial (600 IU orange cap / 1200 IU white cap)
- 2 Prefilled syringe / Prefilled syringes (1 syringe in the 600 IU packaging / 2 syringes in the 1200 IU packaging)
- 3 Needle for mixing the drug
- 4 Disposable syringes for injection (9 syringes in the 600 IU packaging / 18 syringes in the 1200 IU packaging)




Important!

Menopur should be stored in the refrigerator before mixing. The Menopur bottle contains medication for several days of treatment.

Preparing the solution for injection


- 1 Remove the protective cap from the Menopur bottle.
- 2 Pull the rubber cap off the mixing syringe.
- 3 Twist off the protective sleeve from the mixing needle.
- 4 Attach the mixing needle to the mixing syringe.
- 5 Pull the needle cover off the mixing needle.



Mixing the Menopur solution:

- 1 Insert the needle of the mixing syringe through the center mark of the rubber membrane on the vial and slowly inject the liquid into the vial.


The powder dissolves quickly, usually after a few drops of liquid have been transferred to the bottle. Note that all liquid should be transferred.



Note:

If you have been prescribed **Menopur 600 IU**, proceed to point 2.
If you have been prescribed **Menopur 1200 IU**, continue below.

Make sure the needle tip is above the liquid level in the Menopur bottle. Then unscrew the first mixing syringe from the needle. Leave the needle in the **Menopur 1200 IU** bottle.




Take the second mixing syringe that came in the package. **Perform step 2 under the heading “Preparing the solution for injection”** with this syringe as well. Then attach the syringe to the mixing needle that is in the bottle and slowly inject the liquid.

- 2 Then remove the mixing syringe with needle from the Menopur bottle.
- 3 Gently swirl the Menopur bottle to dissolve the powder, do not shake.
- 4 The medicine is now ready to use or store at room temperature ($\leq 25^{\circ}\text{C}$) for up to 28 days.




Injection:

- 1 Take a disposable syringe for injection and pull off the needle cover. Place the Menopur bottle on a firm surface. Insert the needle through the center mark on the rubber membrane on the bottle. Push the plunger into the bottom of the syringe.
- 2 Turn the Menopur bottle and the syringe as shown in the picture and withdraw the desired Menopur dose, see the graduation on the syringe.



Disinfect and inject as instructed by your doctor/nurse.

- 3 Grasp the skin on the right or left side with one hand. Insert the needle into the skin at a 90° angle. Slowly push the plunger all the way in. When the syringe is empty, pull out the needle.
- 4 Vary the injection site as directed by your doctor/nurse.
- 5 Place used needles in a waste container, such as an empty milk carton.
- 6 The remaining medicine in the Menopur bottle can be stored at room temperature ($\leq 25^{\circ}\text{C}$) for up to 28 days.



Note:

The Menopur bottle contains medication for several days of treatment. Therefore, only withdraw the exact dose prescribed by your doctor.





Important!

Before mixing:

Menopur should be stored in a refrigerator.

After mixing:

Menopur can be stored at room temperature ($\leq 25^{\circ}\text{C}$) up to 28 days.

MENOPUR® 600 IU and 1200 IU – (menotropin (human menopausal gonadotropin, HMG)).

Pharmaceutical form: Powder and solvent for solution for injection.

Menopur is indicated for the treatment of infertility in the following clinical situations: Women who cannot become pregnant because their ovaries do not produce eggs (including polycystic ovary syndrome) and who have not responded to treatment with clomiphene citrate and to stimulations the ovaries to develop multiple follicles for assisted reproductive technology (ART) such as in vitro fertilization/embryo transfer (IVF/ET), gamete intra-fallopian transfer (GIFT) and intracytoplasmic sperm injection (ICSI).

Should not be used in women with tumors of the pituitary gland or hypothalamus, ovarian, uterine or breast cancer, gynecological bleeding of unknown cause, ovarian cysts or ovarian enlargement not due to polycystic ovary syndrome, genital malformations or uterine fibroids, or in women who have reached early menopause or during pregnancy and lactation. Treatment requires regular monitoring of ovarian response. Talk to your doctor if you experience stomach pain, abdominal swelling, nausea, vomiting, diarrhea, weight gain, breathing problems or decreased urine production. Treatment with hormones such as Menopur may increase the risk of: multiple pregnancy, miscarriage, ectopic pregnancy, tumors of the reproductive organs, congenital malformations, venous thromboembolism. Read the package leaflet in the medicine package carefully. The medicine is a prescription drug. Always consult a doctor before using the medicine. Benefit limited: Menopur is not subsidized for egg freezing for social reasons. Product summary and package leaflet updated 2018-03-21. For further information and information on prices, see www.fass.se. Ferring Läkemedel AB, Box 4041, 203 11 MALMÖ, Tel: 040-6916900, www.ferring.se

